

Polarization scattering by soliton-soliton collisions

Valery Shchesnovich

Division for Optical Problems in Information Technologies, Academy of Sciences of Belarus, Zhodinskaya Str. 1/2, 220141 Minsk, Republic of Belarus.

Abstract. Collision of two solitons of the Manakov system is studied analytically. Existence of a complete polarization mode switching regime is proved and the parameters of solitons prepared for polarization switching are found.

The idea of an analytical approach to the description of polarized (i.e., vector) solitons in nonlinear optical fibers is based on the fact [1] that for some values of the parameters the original system of coupled nonlinear Schrödinger equations transforms to the Manakov system

$$\begin{aligned} iq_{1z} + \frac{1}{2}q_{1\tau\tau} + (|q_1|^2 + |q_2|^2)q_1 &= 0, \\ iq_{2z} + \frac{1}{2}q_{2\tau\tau} + (|q_2|^2 + |q_1|^2)q_2 &= 0. \end{aligned} \tag{1}$$

This system was shown to be integrable by Manakov [2]. Here q_1 and q_2 are normalized envelopes of the two modes of the soliton, while τ and z are, correspondingly, the normalized retarded time and distance along the fiber. The one-soliton solution for Eq. (1) reads

$$q_l = -2i\eta\theta_l \frac{\exp(-2i[\xi\tau + 2(\xi^2 - \eta^2)z])}{\cosh(2\eta[\tau - \tau_0 + 2\xi z])}, \quad l = 1, 2, \tag{2}$$

where $\theta_l = |\theta_l|e^{i\phi_l}$, $l = 1, 2$, $|\theta_l|$ and ϕ_l are the intensity and initial phase of the l -th polarization component of the soliton, $|\theta_1|^2 + |\theta_2|^2 = 1$, and 2ξ and 2η are the velocity and amplitude of the soliton, correspondingly. Many-soliton solutions of the Manakov system can be easily retrieved with the help of the Riemann-Hilbert problem approach. A detailed analysis of the associated Riemann-Hilbert problem and the perturbation theory for Manakov solitons can be found in Ref. [3]. The general soliton matrices¹, which allow for the

¹Discovered quite recently.

complete classification of all possible soliton solutions to the Manakov system as well, can be found in Ref [4].

Below we will give the formulas describing collision of two Manakov solitons with arbitrary polarization states. To conveniently describe the phase and polarization state of the two-component soliton let us introduce a two-dimensional column vector $|\theta\rangle$, which will be referred to below as the theta-parameter. The Hermitian-transposed row vector will be denoted as $\langle\theta|$. Analysis of the two-soliton solution of the Manakov system (1) under the condition that $\xi_2 > \xi_1$ leads to the following transformation formulas for the parameters $|\theta^{(1)}\rangle$ and $|\theta^{(2)}\rangle$:

$$|\theta^{(+)(1)}\rangle = \exp\left(i\arg\left(\frac{k_1 - \bar{k}_2}{k_1 - k_2}\right) - \beta\right) \left(I + \frac{k_2 - \bar{k}_2}{k_1 - k_2} |\theta^{(-)(2)}\rangle\langle\theta^{(-)(2)}| \right) |\theta^{(-)(1)}\rangle, \quad (3)$$

$$|\theta^{(+)(2)}\rangle = \exp\left(i\arg\left(\frac{k_1 - \bar{k}_2}{\bar{k}_1 - \bar{k}_2}\right) - \beta\right) \left(I + \frac{k_1 - \bar{k}_1}{\bar{k}_1 - \bar{k}_2} |\theta^{(-)(1)}\rangle\langle\theta^{(-)(1)}| \right) |\theta^{(-)(2)}\rangle,$$

where the top signs, minus and plus, denote the theta-parameters before and after the collision, correspondingly, $k_j = \xi_j + \eta_j$, $j = 1, 2$, and β reads

$$\beta = \frac{1}{2} \ln \left(1 - \frac{(k_2 - \bar{k}_2)(k_1 - \bar{k}_1)}{|k_1 - k_2|^2} \left| \langle\theta^{(-)(1)} | \theta^{(-)(2)}\rangle \right|^2 \right). \quad (4)$$

The transformation given by Eqs. (3) and (4) was first derived by Manakov [2] in an equivalent form.

It is easy to verify that the theta-parameters satisfy the following simple identity

$$\langle\theta^{(+)(1)} | \theta^{(+)(2)}\rangle = \exp(4i \arg(k_1 - k_2)) \langle\theta^{(-)(1)} | \theta^{(-)(2)}\rangle. \quad (5)$$

Additionally to transformation (3)-(4), the interaction of two Manakov solitons leads to the finite shifts in the time of arrival parameters of the two solitons:

$$\Delta\tau_o^{(1)} \equiv \tau_o^{(+)(1)} - \tau_o^{(-)(1)} = \frac{1}{2\eta_1} \left(\ln \left| \frac{k_1 - \bar{k}_2}{k_1 - k_2} \right| + \beta \right), \quad (6)$$

$$\Delta\tau_o^{(2)} \equiv \tau_o^{(+)(2)} - \tau_o^{(-)(2)} = -\frac{1}{2\eta_2} \left(\ln \left| \frac{k_1 - \bar{k}_2}{k_1 - k_2} \right| + \beta \right).$$

Equations (3), (4), and (6) completely describe the collision of two Manakov solitons.

The transformation given by Eqs. (3)-(4) means that the colliding solitons significantly alter each other's polarizations and phases. This fact was already mentioned by Manakov [2]. Moreover, there are the polarization switching regimes of collision, i.e. when we have $\theta_l^{(+)(n)} = 0$ but $\theta_l^{(-)(n)} \neq 0$, for some n and l ($n = 2$ and $l = 2$ below). The polarization switching regime was discovered in Ref. [5] in the numerical simulations of the general two-soliton solution. To analytically prove the existence of such a regime let us invert one of the equations from system (3) for the theta-parameters of the solitons. For instance, given $|\theta^{(+)(2)}\rangle$ and $|\theta^{(-)(1)}\rangle$ we obtain for the rest of the theta-parameters²:

$$|\theta^{(-)(2)}\rangle = \exp\left(i\arg\left(\frac{\bar{k}_1 - k_2}{k_1 - \bar{k}_2}\right) + \beta\right) \left(I + \frac{k_1 - \bar{k}_1}{\bar{k}_2 - k_1} |\theta^{(-)(1)}\rangle\langle\theta^{(-)(1)}| \right) |\theta^{(+)(2)}\rangle, \quad (7)$$

$$|\theta^{(+)(1)}\rangle = \exp\left(i\arg\left(\frac{k_1 - \bar{k}_2}{k_1 - k_2}\right) + \beta\right) \left(I + \frac{k_2 - \bar{k}_2}{\bar{k}_1 - k_2} |\theta^{(+)(2)}\rangle\langle\theta^{(+)(2)}| \right) |\theta^{(-)(1)}\rangle.$$

Importantly, the set of the theta-parameters $\{|\theta^{(-)(1)}\rangle, |\theta^{(+)(2)}\rangle\}$ is a close one due to the fact that β admits also the following equivalent representation

$$\beta = -\frac{1}{2} \ln \left(1 + \frac{(k_2 - \bar{k}_2)(k_1 - \bar{k}_1)}{|k_1 - \bar{k}_2|^2} \left| \langle\theta^{(-)(1)}|\theta^{(+)(2)}\rangle \right|^2 \right). \quad (8)$$

Let us put

$$|\theta^{(-)(1)}\rangle = \begin{pmatrix} \cos\alpha_1 e^{i\phi_1} \\ \sin\alpha_1 e^{i\phi_2} \end{pmatrix}, \quad |\theta^{(+)(2)}\rangle = \begin{pmatrix} e^{i\psi} \\ 0 \end{pmatrix}, \quad (9)$$

where α_1, ϕ_1, ϕ_2 and ψ are given. Define also

$$|\theta^{(-)(2)}\rangle = \begin{pmatrix} \cos\alpha_2 e^{i\psi_1} \\ \sin\alpha_2 e^{i\psi_2} \end{pmatrix}. \quad (10)$$

²Another way to derive Eqs. (7)-(8) is to use the asymptotic expansion of general two-soliton solution on the two characteristics, defined by $\tau + 2\xi_n z = \text{const.}$, $n = 1, 2$, and the fact that there are exactly two equivalent factored representations of the soliton matrix: $\Gamma(k) = \chi_1(k)\chi_2(k) = \widetilde{\chi}_2(k)\widetilde{\chi}_1(k)$, for notations consult Ref. [4].

For the initial (i.e. before the collision) theta-parameter of soliton 2 we obtain from Eq. (7)

$$| \theta^{(-)(2)} \rangle = \frac{\exp \left(i \arg \left(\frac{k_2 - \bar{k}_1}{k_2 - k_1} \right) + i \psi \right)}{\left(1 - \frac{4\eta_1\eta_2}{|k_2 - k_1|^2} \cos^2 \alpha_1 \right)^{\frac{1}{2}}} \begin{pmatrix} 1 + \frac{2i\eta_1}{k_2 - k_1} \cos^2 \alpha_1 \\ \frac{2i\eta_1}{k_2 - k_1} \cos \alpha_1 \sin \alpha_1 e^{(i\phi_2 - i\phi_1)} \end{pmatrix} \quad (11)$$

One can use Eqs. (10) and (11) to find the values of the polarization angle and phase difference of soliton 2 prepared for the complete polarization switching due to collision with soliton 1 with the phase vector fixed by Eq. (9):

$$\alpha_2 = -\arctan \left(\frac{\eta_1 \sin(2\alpha_1)}{\left((\xi_2 - \xi_1)^2 + (\eta_2 - \eta_1 \cos(2\alpha_1))^2 \right)^{\frac{1}{2}}} \right), \quad (12)$$

$$\psi_2 - \psi_1 = \phi_2 - \phi_1 + \frac{\pi}{2} + \arctan \left(\frac{\eta_2 - \eta_1 \cos(2\alpha_1)}{\xi_2 - \xi_1} \right).$$

Therefore, it is analitically shown that vector solitons governed by the system of coupled nonlinear Schrödinger equations exhibit a novel type of soliton collision – the soliton polarization state switching. In the case when the Manakov system is the relevant model the polarization switching will be complete. The soliton polarization state switching can be used for the logic elements based on vector solitons.

Acknowledgement

The work was supported by the Fundamental Research Foundation of Belarus under contract No. MII96-06. The author is grateful to Dr. Takayuki Tsuchida for pointing out a misprint.

References

- [1] C. R. Menyuk, IEEE J. Quant. Electron. **25**, 2674 (1989).
- [2] S. V. Manakov, Zh. Eksp. Teor. Fiz. **65**, 505 (1973) [Sov. Phys. JETP **38**, 248 (1974)].

- [3] V.S. Shchesnovich and E.V. Doktorov, Phys. Rev. E **55**, 7626 (1997).
- [4] V. S. Shchesnovich and J. Yang, J. Math. Phys. **44**, 4604 (2003).
- [5] R. Radhakrishnan, M. Lakshmanan, and J. Hietarinta, Phys. Rev. E. **56**, 2213 (1997).